

# STDs and HIV – CDC Fact Sheet



***People who have STDs are more likely to get HIV, when compared to people who do not have STDs.***



## **Are some STDs associated with HIV?**

Yes. In the United States, people who get syphilis, gonorrhea, and herpes often also have HIV, or are more likely to get HIV in the future.

## **Why does having an STD put me more at risk for getting HIV?**

If you get an STD you are more likely to get HIV than someone who is STD-free. This is because the same behaviors and circumstances that may put you at risk for getting an STD can also put you at greater risk for getting HIV. In addition, having a sore or break in the skin from an STD may allow HIV to more easily enter your body.

## **What activities can put me at risk for both STDs and HIV?**

- Having anal, vaginal, or oral sex without a condom;
- Having multiple sex partners;
- Having anonymous sex partners;
- Having sex while under the influence of drugs or alcohol can lower inhibitions and result in greater sexual risk-taking.

## **What can I do to prevent getting STDs and HIV?**

The only way to avoid STDs is to not have vaginal, anal, or oral sex. If you are sexually active, you can do the following things to lower your chances of getting STDs and HIV:

- Choose less risky sexual behaviors.
- Use condoms consistently and correctly.
- Reduce the number of people with whom you have sex.
- Limit or eliminate drug and alcohol use before and during sex.
- Have an honest and open talk with your healthcare provider and ask whether you should be tested for STDs and HIV.
- Talk to your healthcare provider and find out if pre-exposure prophylaxis, or PrEP, is a good option for you to prevent HIV infection.

